

020026

10/687,478

## COMPLETE LISTING OF CLAIMS

1. (Original) A method of estimating a parameter of a local maxima or minima of a function comprising:
  - performing interpolation on samples of the function at or near a local maxima or minima, resulting in an interpolated local maxima or minima;
  - deriving an interpolation offset comprising a deviation between locations of the interpolated local maxima or minima and a sampled local maxima or minima; and
  - deriving an estimate of the parameter from the interpolation offset.
2. (Original) The method of claim 1 wherein the function is a correlation function.
3. (Original) The method of claim 2 wherein the correlation function is derived from a received signal.
4. (Original) The method of claim 2 wherein the second deriving step comprises deriving a parameter bias from the interpolation offset using a pre-existing relationship that is present between these two variables and then deriving an estimate of the parameter from the parameter bias.
5. (Original) The method of claim 4 wherein the interpolation offset comprises a deviation between locations of interpolated and sampled peaks along a code phase dimension.
6. (Original) The method of claim 5 wherein the parameter bias is a code phase bias.
7. (Original) The method of claim 6 wherein the parameter which is estimated is location of a peak along the code phase dimension, and an estimate of this parameter is derived from the code phase bias.

BEST AVAILABLE COPY

020026

10/687,478

8. (Original) The method of claim 4 wherein the interpolation offset comprises a deviation between locations of interpolated and sampled peaks along a Doppler frequency dimension.

9. (Original) The method of claim 8 wherein the parameter bias is a Doppler frequency bias.

10. (Original) The method of claim 9 wherein the parameter which is estimated is location of a peak of the function along the Doppler frequency dimension, and an estimate of this parameter is derived from the Doppler frequency bias.

11. (Previously Presented) The method of claim 4 wherein the parameter bias is a peak energy bias.

12. (Original) The method of claim 11 wherein the parameter which is estimated is peak energy, and an estimate of this parameter is derived from the peak energy bias.

13. (Cancelled)

14. (Cancelled)

15. (Original) The method of claim 4 wherein the pre-existing relationship between the interpolation offset and the parameter bias is embodied as a lookup table.

16. (Original) The method of claim 15 wherein the second deriving step comprises directly deriving an estimate of the parameter from the interpolation offset through an access to the lookup table.

17. (Withdrawn) A memory tangibly embodying a lookup table, the lookup table implementing a pre-existing relationship between an interpolation offset and a parameter bias or parameter estimate, and the interpolation offset comprising a deviation between locations of interpolated and sampled local maxima or minima of a function.

020026

10/687,478

18. (Withdrawn) The memory of claim 17 wherein the function is a correlation function.

19. (Withdrawn) The memory of claim 18 wherein the correlation function is derived from a received signal.

20. (Withdrawn) The memory of claim 18 wherein an access to the lookup table yields the parameter bias.

21. (Withdrawn) The memory of claim 18 wherein an access to the lookup table yields the parameter estimate.

22. (Withdrawn) The memory of claim 20 wherein the interpolation offset comprises a deviation between locations of interpolated and sampled peaks along a code phase dimension.

23. (Withdrawn) The memory of claim 22 wherein the parameter bias is a code phase bias.

24. (Withdrawn) The memory of claim 23 wherein the parameter estimate is an estimate of the location of a peak along a code phase dimension.

25. (Cancelled)

26. (Withdrawn) The memory of claim 17 wherein the parameter bias is a peak energy bias.

27. (Withdrawn) The memory of claim 26 wherein the parameter estimate is an estimate of peak energy.

28. (Withdrawn) The memory of claim 26 wherein the parameter estimate comprises a sum of interpolated peak energy and the peak energy bias.

020026

10/687,478

29. (Withdrawn) The memory of claim 26 wherein the parameter estimate comprises a sum of sampled peak energy and the peak energy bias.

30. (Withdrawn) The memory of claim 17 wherein the interpolation offset comprises a deviation between locations of interpolated and sampled peaks along a Doppler frequency dimension.

31. (Withdrawn) The memory of claim 30 wherein the parameter bias is a Doppler frequency bias.

32. (Withdrawn) The memory of claim 31 wherein the parameter estimate is an estimate of the location of a peak along the Doppler frequency dimension.

33. (Withdrawn) The memory of claim 31 wherein the parameter estimate comprises a sum of the interpolation offset and the Doppler frequency bias.

34. (Cancelled)

35. (Cancelled)

36. (Cancelled)

37. (Cancelled)

38. (Withdrawn) The memory of claim 17 in combination with a processor, wherein the processor is configured to access the lookup table.

39. (Original) A memory tangibly embodying a sequence of software instructions for performing a method of estimating a parameter of a local maxima or minima of a function comprising:

020026

10/687,478

performing interpolation on samples of the function at or near a local maxima or minima, resulting in an interpolated local maxima or minima;

deriving an interpolation offset comprising a deviation between locations of the interpolated local maxima or minima and a sampled local maxima or minima; and

deriving an estimate of the parameter from the interpolation offset.

40. (Original) The memory of claim 39 wherein the function is a correlation function.

41. (Original) The memory of claim 40 wherein the correlation function is derived from a received signal.

42. (Previously Presented) The memory of claim 40 wherein deriving an estimate of the parameter from the interpolation offset comprises deriving a parameter bias from the interpolation offset using a pre-existing relationship which is present between these two variables and deriving an estimate of the parameter from the parameter bias.

43. (Original) The memory of claim 42 wherein the interpolation offset comprises a deviation between locations of interpolated and sampled peaks along a code phase dimension.

44. (Original) The memory of claim 43 wherein the parameter bias is a code phase bias.

45. (Original) The memory of claim 44 wherein the parameter is location of a peak along the code phase dimension, and an estimate of this parameter is derived from the code phase bias.

46. (Original) The memory of claim 42 wherein the interpolation offset comprises a deviation between locations of interpolated and sampled peaks along a Doppler frequency dimension.

47. (Original) The memory of claim 46 wherein the parameter bias is a Doppler frequency bias.

020026

10/687,478

48. (Original) The memory of claim 47 wherein the parameter which is estimated is location of a peak of the function along the Doppler frequency dimension, and an estimate of this parameter is derived from the Doppler frequency bias.

49. (Previously Presented) The memory of claim 42 wherein the parameter bias is a peak energy bias.

50. (Original) The memory of claim 49 wherein the parameter which is estimated is peak energy, and an estimate of this parameter is derived from the peak energy bias.

51. (Original) The memory of claim 46 wherein the parameter bias is a peak energy bias.

52. (Original) The memory of claim 51 wherein the parameter which is estimated is peak energy, and an estimate of this parameter is derived from the peak energy bias.

53. (Original) The memory of claim 42 wherein the pre-existing relationship between the interpolation offset and the parameter bias is embodied as a lookup table.

54. (Previously Presented) The memory of claim 39 in combination with a processor, wherein the processor is configured to access and execute the sequence of software instructions tangibly embodied by the memory.

55. (Previously Presented) An apparatus for estimating a parameter of a local maxima or minima of a function comprising:

means for performing interpolation on samples of the function at or near a local maxima or minima, resulting in an interpolated local maxima or minima;

means for deriving an interpolation offset comprising a deviation between locations of the interpolated local maxima or minima and a sampled local maxima or minima; and

means for deriving an estimate of the parameter from the interpolation offset.

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☒ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**